

Utility maximization with current utility on the wealth: regularity of solutions to the HJB equation *

Salvatore Federico^{a)}Paul Gassiat^{b)}Fausto Gozzi^{c)}

December 11, 2013

Abstract

We consider a utility maximization problem for an investment-consumption portfolio when the current utility depends also on the wealth process. Such kind of problems arise, e.g., in portfolio optimization with random horizon or with random trading times. To overcome the difficulties of the problem we use the dual approach. We define a dual problem and treat it by means of dynamic programming, showing that the viscosity solutions of the associated Hamilton-Jacobi-Bellman equation belong to a suitable class of smooth functions. This allows to define a smooth solution of the primal Hamilton-Jacobi-Bellman equation, proving that this solution is indeed unique in a suitable class and coincides with the value function of the primal problem. Some financial applications of the results are provided.

Keywords: Optimal stochastic control, investment-consumption problem, duality, Hamilton-Jacobi-Bellman equation, regularity of viscosity solutions.

MSC 2010 Classification: 93E20, 49L20, 90C46, 91G80, 35B65.

JEL Classification: C61, G11.

Contents

1	Introduction	2
2	Model and optimal control problem	3
3	Primal and dual control problem	7

*This research has been partially supported by the PRIN research project “Metodi deterministici e stocastici nello studio di problemi di evoluzione” of the Italian Minister of University and Research.

^{a)} Dipartimento di Economia, Management e Metodi Quantitativi, Università di Milano, Italy. E-mail: salvatore.federico@unimi.it

^{b)} Institut für Mathematik, TU Berlin. E-mail: gassiat@math.tu-berlin.de

^{c)} Dipartimento di Economia e Finanza, Libera Università degli Studi Sociali “Guido Carli”, Roma, Italy. E-mail: fgozzi@luiss.it

4	The dual value function as classical solution of the dual HJB equation	11
4.1	W as viscosity solution of the dual HJB equation	11
4.2	Regularity of W	17
5	Back to the primal control problem: verification and optimal controls	19
5.1	\widetilde{W} is a classical solution of the primal HJB equation	20
5.2	Verification theorem	21
5.3	Optimal feedback controls	23
6	Applications	25
6.1	Portfolio optimization with random horizon	25
6.2	Market with liquid and illiquid assets	26

1 Introduction

The paper deals with the problem of utility maximization in consumption-investment models over a fixed horizon when the current utility depends also on the wealth process. The fact that the current utility may depend also on the wealth is motivated by the fact that this situation arises in some concrete financial problems, as discussed in Section 6.

We tackle the problem by duality and using a dynamic programming approach both on the primal and on the dual problem. Since the papers by Karatzas, Lehoczky and Shreve [19] and by Cox and Huang [7], the duality approach to consumption-investment problems has been extensively treated in the literature (see the survey paper by Rogers [24], and the book by Karatzas and Shreve [20, Ch. 3 and 6] - and the references therein) to treat generalizations of the classical Merton problem (incomplete markets, non-Markovian setting, strategies constraints, transaction costs, etc.). Notably with regard to our paper, Bouchard and Pham [4] treat the case of current utility depending on the wealth in a semimartingale setting without developing the dynamic programming approach.

When the stock is assumed to evolve according to a stochastic differential equation, one can apply the dynamic programming machinery both to the primal and the dual problem to get some more insights on the solution of the problem. In particular the duality can be read at the analytical level of the Hamilton-Jacobi-Bellman (HJB) equation, providing a dual equation. This is what is done in Bian, Miao and Zheng [2] (see also the extension of such results in [3]) in the case of no current utility on the wealth. But, as far as we know, duality has been never employed combined with the dynamic programming when the current utility depends on the wealth process. This may be due to the fact that when there is no dependence of the current utility on the wealth process the HJB equation associated to the dual problem is linear - so approachable by semi-explicit solution written in terms of the heat kernel (see [2, 3]) - while when the current utility also depends on the wealth such HJB equation is just semi-linear - so more difficult to study. At the level of control problems, this corresponds to the fact that in the former case the dual problem is simpler, as the control does not appear in it, while in the latter one the dual problem is a real control problem (these issues are discussed in Remark 3.1). Nevertheless, also in

this last case, the dual control problem is still simpler to treat than the primal one, as the control only appears in the drift of the process, consistently with the fact that the HJB equation is semilinear (while the HJB equation associated to the primal control problem is fully nonlinear and degenerate, so very difficult to tackle directly by the PDE's theory of classical solutions).¹

Our method to solve the problem is the following.

Step 1: Starting from the original primal problem (with value function V and an associated primal HJB equation), we define a dual problem, which is still a control problem,

Step 2: We associate to the dual problem a dual HJB equation and prove that the value function W of this dual problem is a viscosity solution of the dual HJB equation (Proposition 4.4).

Step 3: Since the dual HJB equation is semilinear and nondegenerate, we are able to prove good regularity results for W . This is proved in Theorem 4.5, which is the key result of the paper.

Step 4: The regularity W allows to define a smooth solution to the primal HJB equation, which is the Legendre transform \widetilde{W} of W .

Step 5: We prove a verification theorem for our primal problem within a suitable class \mathcal{C} of smooth solutions of the primal HJB equation. Since $\widetilde{W} \in \mathcal{C}$, this theorem, together with a result of existence and uniqueness for the associated closed loop equation, will imply that $\widetilde{W} = V$ and that V is the unique classical solution of the primal HJB equation within the class \mathcal{C} . These results will yield also the construction of an optimal feedback control for the primal problem.

The rest of the paper is organized as follows. In Section 2 we set the problem and state the assumptions. In Section 3 we define the dual problem (Step 1 above). In Section 4 we study the dual HJB equation by a viscosity approach and state the regularity of the value function W (Steps 2 and 3 above). In Section 5 we prove that V is a classical solution of the HJB equation and provide the optimal feedbacks through a verification theorem (Steps 4 and 5 above). Finally, Section 6 provides two concrete applications of our framework.

2 Model and optimal control problem

In this section we present the financial model and the (primal) stochastic control problem we deal with.

Let us consider a complete filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ satisfying the usual conditions, on which is defined a standard Brownian motion $(B_t)_{t \geq 0}$. We assume that $(\mathcal{F}_t)_{t \geq 0}$ is the filtration generated by this Brownian motion and enlarged by the \mathbb{P} -null sets.

On this space we consider a riskless asset with deterministic rate of return that without loss of generality (see Remark 2.3(ii) below) we set equal to 0, and a risky asset $S = (S_t)_{t \geq 0}$ with dynamics

$$dS_t = S_t(b(t)dt + \sigma(t)dB_t),$$

¹We also should mention the paper [23], where the HJB equation associated to the dual problem is again fully nonlinear, but admits a semi-explicit solution in the form of a power series.

where b, σ are deterministic coefficients representing, respectively, the drift and the volatility of the risky asset.

Fix a time horizon $T > 0$. In the setting above, we define a set of admissible trading/consumption strategies in the following way. Consider all the couples of processes (c, π) such that

- (h1) $c = (c_t)_{t \in [0, T]}$ is a real nonnegative process $(\mathcal{F}_t)_{t \in [0, T]}$ -predictable and with trajectories locally integrable in $[0, T]$; c_t represents the consumption rate at time t ;
- (h2) $\pi = (\pi_t)_{t \in [0, T]}$ is a real process $(\mathcal{F}_t)_{t \in [0, T]}$ -predictable and with trajectories locally square integrable in $[0, T]$; π_t represents the amount of money invested in the risky asset at time t .

Given a couple (c, π) satisfying the requirements (h1)-(h2) above, we can consider the process X_t representing the wealth associated to such strategy. Its dynamics is

$$\begin{cases} dX_t = \pi_t(b(t)dt + \sigma(t)dB_t) - c_t dt, \\ X_0 = x_0, \end{cases} \quad (1)$$

where $x_0 \geq 0$ is the initial wealth. As class of admissible controls we consider the couples of processes (c, π) satisfying (h1)-(h2) and such that the corresponding wealth process X is nonnegative (no-bankruptcy constraint). The optimization problem is

$$\mathbb{E} \left[\int_0^T U_1(t, c_t, X_t) dt + U_2(X_T) \right]. \quad (2)$$

Let $\mathbb{R}_+ := [0, +\infty)$. In the following, given an integer $k \geq 0$, a real number $\delta > 0$ and $\mathcal{O} \subset \mathbb{R}^n$ open, by $C_{loc}^{\delta/2, k+\delta}([0, T] \times \mathcal{O}; \mathbb{R})$ we denote the space of real continuous functions on $[0, T] \times \mathcal{O}$ whose space derivatives up to order k exist and are $\delta/2$ -Hölder continuous with respect to t and δ -Hölder continuous with respect to the space variables on each compact subset of $[0, T] \times \mathcal{O}$. We make the following assumptions on the model.

Assumption 2.1 $b, \sigma : [0, T] \rightarrow \mathbb{R}$ are strictly positive and $(\delta/2)$ -Hölder continuous for some $\delta \in (0, 1]$.

Assumption 2.2 The preference of the agent are described by utility functions U_1, U_2 satisfying the following:

- (i) $U_1 : [0, T] \times \mathbb{R}_+^2 \rightarrow \mathbb{R}$ is such that $U_1 \in C_{loc}^{\delta/2, k+\delta}([0, T] \times (0, +\infty) \times (0, +\infty); \mathbb{R})$ for some $k \geq 2$ (and the same δ of Assumption 2.1). For each fixed $t \in [0, T]$ the function $U_1(t, \cdot, \cdot)$ is concave with respect to (c, x) and nondecreasing with respect to both the variables c, x .

Moreover either

$$(a) \quad \begin{cases} \frac{\partial}{\partial c} U_1(t, 0^+, x) = +\infty, & \forall (t, x) \in [0, T) \times \mathbb{R}_+, \\ \frac{\partial}{\partial c} U_1(t, +\infty, x) = 0, & \forall (t, x) \in [0, T) \times \mathbb{R}_+, \\ \frac{\partial}{\partial c} U_1 > 0, \quad \frac{\partial^2}{\partial c^2} U_1 < 0, & \text{in } [0, T) \times (0, +\infty) \times (0, +\infty), \end{cases}$$

or

$$(b) \quad \frac{\partial}{\partial c} U_1 \equiv 0.$$

(ii) $U_2 : \mathbb{R}_+ \rightarrow \mathbb{R}$ is continuous, nondecreasing, concave. Without loss of generality we assume

$$U_2(0) = 0. \quad (3)$$

(iii) The following growth condition holds: there exist $K > 0$ and $p \in (0, 1)$ such that

$$U_1(t, c, x) + U_2(x) \leq K(1 + c^p + x^p), \quad \forall (t, c, x) \in [0, T) \times \mathbb{R}_+^2. \quad (4)$$

Moreover, without loss of generality for the optimization problem, we assume that

$$U_1(t, 0, 0) = 0, \quad \forall t \in [0, T). \quad (5)$$

(iv) Either

$$(a) \quad \exists \varepsilon > 0 \text{ such that } \lim_{c \rightarrow +\infty} U_1(t, c, 0) = +\infty \text{ uniformly in } t \in [T - \varepsilon, T),$$

or

$$(b) \quad \lim_{x \rightarrow +\infty} U_2(x) = +\infty$$

or both.

In the remark below we comment on some features of the model and explain when and how they can be eventually modified to cover other interesting cases.

Remark 2.3 (i) In the applications one is often interested to work with power utility functions. Assumption 2.2 includes only the case of positive power. On one hand the case of negative exponent is interesting, as it seems to be even more realistic from the point of view of the agents' behavior; on the other hand, it would require a slightly different treatment. Just for simplicity, we will work with Assumption 2.2, nevertheless we stress that the case of negative utility can be treated by the same techniques by suitable modifications.

(ii) The assumption that the riskless rate of return is 0 can be done without loss of generality. Indeed, since we are considering a quite general time-dependent U_1 , the interest rate can be discarded in it by a suitable discounting of the variables (see [17, Rem. 2, p. 189]).

(iii) The problem without consumption falls in our setting as well. Indeed, take a problem without consumption and with running utility $u_1(t, x)$. Defining $U_1(t, c, x) = u_1(t, x)$ in

our setting, consuming turns out to be not convenient, as its negative effect on the wealth has not a trade-off in terms of utility from consumption. In other terms, the optimal consumption is $c_t^* \equiv 0$. As a consequence the problem in our setting with U_1 defined as above is equivalent to the problem without consumption and with utility function u_1 . In particular, when $u_1 \equiv 0$ we fall in the setting of [2].

(iv) We have set the problem with finite horizon. However, some problems arising in the applications - see Section 6 - involve the infinite horizon case, where $T = +\infty$, for which the functional usually looks like

$$\mathbb{E} \left[\int_0^\infty e^{-\rho t} U_1(t, c_t, X_t) dt \right],$$

where, as usual for infinite horizon problems, $\rho > 0$ is a discount rate sufficiently large to guarantee the finiteness of the value function. The results we provide in the present paper for the finite horizon case can be suitably generalized to the infinite horizon case, with the complication of dealing in the viscosity treatment of the HJB equation with growth conditions for $t \rightarrow +\infty$ in place of terminal boundary conditions at $t = T$. We refer, e.g., to [12] for an example of the technical treatment of this kind of conditions and stress here that our main results - the regularity results - do not “see” whether the horizon is finite or infinite, as they are based on local arguments. Of course, in this case one needs to assume that Assumption 2.2(iv) is satisfied at point (a).

(v) We comment on Assumption 2.2(i). It requires that either U_1 is independent of c or it satisfies Inada’s conditions with respect to c . We need this assumption to get in a straightforward way the regularity of the Legendre transform of U_1 with respect to c (Proposition 4.1(6)), which is in turn needed to get the regularity of the dual value function, see Section 4.2. Basically it is thought to cover the case of separable utility in the form $U_1(t, c, x) = U_1^{(1)}(t, c) + U_1^{(2)}(t, x)$, where $U_1^{(1)}$ is identically 0 or satisfies the Inada conditions with respect to c , which is the case arising in the applications we have in mind (see Section 6). Relaxing this assumption seems possible, but at a price of more demanding technical arguments. We prefer to avoid such technicalities in order to focus on the main topic of the paper, which is the regularity of solutions of the HJB equation by means of the duality approach.

(vi) The assumption of strict positivity of b, σ is done to have strict parabolicity of the HJB equation. Actually this is needed only in the interior, so we might allow the cases $b(T) = 0$ and/or $\sigma(T) = 0$. However, allowing that would bring some other technicalities, so we prefer to impose strict positivity also at T . We also stress that we actually need just the assumption $b(t) \neq 0$ for all $t \in [0, T]$; but, due to continuity, this is equivalent to say that b keeps the sign. Since the assumption making sense from a financial point of view is $b(\cdot) > 0$, we impose it.

(vii) Although for simplicity we consider in our model the case of just one risky asset, it is easy to see that the program we described in the introduction works also in more dimensions (more risky assets, as in [2]). In that case strict positivity of $b(t)$ and $\sigma(t)$ in Assumption 2.1 should be replaced by the assumption that for all $t \in [0, T]$ (the matrix) $\sigma(t)$ is invertible and (the vector) $b(t) \neq 0$, so that in the dual HJB equation (25) the term $|\sigma^{-1}(t)b(t)|^2$ is then still well-defined and strictly positive.

(viii) We are concerned with a utility maximization problem. Nevertheless, our approach seems applicable also to different cases, e.g. to the case of quadratic risk minimization, by suitably adapting the arguments.

3 Primal and dual control problem

Since we are going to apply the dynamic programming techniques, we define the optimization problem for generic initial data $(t, x) \in [0, T] \times \mathbb{R}_+$. Let $t \in [0, T]$ and consider all the couples of processes (c, π) such that

(h1') $c = (c_s)_{s \in [t, T]}$ is a real nonnegative process $(\mathcal{F}_s)_{s \in [t, T]}$ -predictable and with trajectories locally integrable in $[0, T]$.

(h2') $\pi = (\pi_s)_{s \in [t, T]}$ is a real process $(\mathcal{F}_s)_{s \in [t, T]}$ -predictable and with trajectories square locally integrable in $[0, T]$.

Given $x \geq 0$ and a couple (c, π) satisfying the requirements (h1')-(h2') above, we denote by $X^{t, x, c, \pi}$ the solution to (1) starting at time t from x and under the control (c, π) . We define a class of admissible controls $\mathcal{A}(t, x)$ depending on the initial $(t, x) \in [0, T] \times [0, +\infty)$ as the set of couples (c, π) satisfying the requirement above and such that the corresponding state trajectory $X^{t, x, c, \pi}$ is nonnegative. We notice that such set is nonempty for each $t \in [0, T]$ and $x \geq 0$, as for such initial data the null strategy $(c, \pi) \equiv (0, 0)$ is always admissible. Moreover $\mathcal{A}(t, x) = \{(0, 0)\}$ if and only if $x = 0$. Then we define the functional

$$J(t, x; c, \pi) := \mathbb{E} \left[\int_t^T U_1(s, c_s, X_s^{t, x, c, \pi}) dt + U_2(X_T^{t, x, c, \pi}) \right].$$

We call *primal control problem* - and denote it by **(P)** - the optimization problem

$$\textbf{(P)} \quad \sup_{(c, \pi) \in \mathcal{A}(t, x)} J(t, x; c, \pi),$$

and denote by V the value function associated to this problem - that we call *primal value function*, i.e.

$$V(t, x) := \sup_{(c, \pi) \in \mathcal{A}(t, x)} J(t, x; c, \pi), \quad (t, x) \in [0, T] \times \mathbb{R}_+.$$

Due to the fact that the state 0 is an absorbing boundary for the problem and to (5)-(3), we see that V satisfies the boundary condition

$$V(t, 0) = 0, \quad \forall t \in [0, T]. \quad (6)$$

On the other hand V clearly satisfies also the terminal condition

$$V(T, x) = U_2(x). \quad (7)$$

Set

$$D_T := [0, T] \times (0, +\infty).$$

By standard arguments of stochastic control (see e.g. [25, Ch.4]), we can associate to V a HJB equation in D_T , which we call *primal* HJB equation. It is

$$-v_t(t, x) - \sup_{c \geq 0, \pi \in \mathbb{R}} H_{cv}(t, x, v_x(t, x), v_{xx}(t, x); c, \pi) = 0, \quad (8)$$

where the function H_{cv} is defined for $(t, x, y, Q) \in D_T \times \mathbb{R}^2$, $c \geq 0, \pi \in \mathbb{R}$, as

$$H_{cv}(t, x, y, Q; c, \pi) := U_1(t, c, x) + (b(t)\pi - c)y + \frac{\sigma(t)^2}{2}\pi^2 Q.$$

When $y > 0$ and $Q < 0$ (which is the case we will consider), the Hamiltonian

$$H(t, x, y, Q) := \sup_{c \geq 0, \pi \in \mathbb{R}} H_{cv}(t, x, y, Q; c, \pi)$$

is finite and takes the form

$$H(t, x, y, Q) = U_1^*(t, y, x) - \frac{b^2(t)}{2\sigma^2(t)} \frac{y^2}{Q}, \quad (9)$$

where U_1^* is the sup-Legendre transform of U_1 with respect to c , i.e. the function (convex in y)

$$U_1^*(t, y, x) := \sup_{c \geq 0} \{U_1(t, c, x) - cy\}, \quad (t, y, x) \in [0, T) \times (0, +\infty) \times \mathbb{R}_+.$$

We expect that V may be characterized as solution of (8) completed by the boundary and terminal conditions (6)-(7). We do not tackle directly the above equation (8), even if a characterization of V as unique viscosity solution to it could be performed.² We just note here that this equation is fully nonlinear and degenerate, so the regularity of its solutions cannot be obtained dealing directly with it by the known methods of classical solutions of PDE's.³ What we can do is to apply duality to the problem and get a dual control problem with an associated HJB equation for which we are able to prove regularity results. For this purpose, consider, for $(t, y) \in [0, T) \times (0, +\infty)$, the sup-Legendre transform of $U_1^*(t, y, \cdot)$, i.e. the function (convex in (y, u))

$$\begin{aligned} \widetilde{U}_1^*(t, y, u) &:= \sup_{x \geq 0} \{U_1^*(t, y, x) - xu\}, \\ &= \sup_{c, x \geq 0} \{U_1(t, c, x) - cy - xu\}, \quad (t, y, u) \in [0, T) \times (0, +\infty) \times (0, +\infty). \end{aligned} \quad (10)$$

²One could try to prove the continuity of V , then show that V is a viscosity solution of the HJB equation and finally use quite standard analytical techniques to prove a comparison in the viscosity sense for the equation and therefore get uniqueness for it (see e.g. [6, 14, 25]). Otherwise one could try to drop the proof of the continuity and deal with discontinuous viscosity solutions, for which the comparison is a bit harder to prove (see [14, Ch. VII]), and then prove the continuity a posteriori as a consequence of the characterization as viscosity solution. We will not do that, since our study of the dual HJB equation will be sufficient to come back and prove a characterization of V as classical solution to the HJB equation within a suitable class of smooth functions. Our uniqueness result will be weaker than what can be obtained by the viscosity approach, but will be enough for our purposes.

³To this regard, we should mention, e.g., [5, 8, 26] for direct results in this direction, when the problem is autonomous and over an infinite horizon, and the equation elliptic. Up to our knowledge, despite a sketch in [26], there are no results of this kind for parabolic HJB equations coming from investment-consumption problems - as the one we deal with in this paper.

For convenience of the reader, we notice that, when U_1 is separable in x and c , i.e. $U_1(t, c, x) = U_1^{(1)}(t, c) + U_1^{(2)}(t, x)$, we have

$$\widetilde{U}_1^*(t, y, u) = \widetilde{U}_1^{(1)}(t, c) + \widetilde{U}_1^{(2)}(t, x),$$

where $\widetilde{U}_1^{(1)}, \widetilde{U}_1^{(2)}$ are, respectively, the sup-Legendre transform of $U_1^{(1)}, U_1^{(2)}$ with respect to the second variable. Finally, we consider also the sup-Legendre transform of U_2 , i.e. the function

$$\widetilde{U}_2(y) = \sup_{x \geq 0} \{U_2(x) - xy\}, \quad y > 0. \quad (11)$$

Given $(t, y) \in \overline{D_T}$, we may consider a new control problem - which we call *dual control problem* and denote by **(D)** - that we are going to define (for the derivation of the argument see [24, Sec. 1]). Let $\beta = (\beta_s)_{s \in [t, T]}$ be a fixed adapted process with locally bounded integrable trajectories and consider the controlled process $Y^{t, y, \beta, u}$ defined by the SDE

$$\begin{cases} dY_s = -u_s ds + \beta_s Y_s dW_s, \\ Y_t = y, \end{cases} \quad (12)$$

with $u \in \mathcal{U}_\beta(t, y)$, where

$$\mathcal{U}_\beta(t, y) = \{(u_s)_{s \in [t, T]} \text{ is } (\mathcal{F}_s)_{s \in [t, T]} \text{-predictable, nonnegative, with integrable trajectories, and such that } Y_s^{t, y, \beta, u} > 0 \text{ a.s. } \forall s \in [t, T]\}. \quad (13)$$

Let $x \in \mathbb{R}_+$, $y > 0$ (c, π) $\in \mathcal{A}(t, x)$, $u \in \mathcal{U}_\beta(t, y)$, and set $X = X^{t, x, c, \pi}$ and $Y = Y^{t, y, \beta, u}$. Integration by parts yields

$$d(X_s Y_s) = (-u_s X_s - c_s Y_s) ds + Y_s (\pi_s \sigma(s) + \beta_s X_s) dW_s + Y_s \pi_s (b(s) + \beta_s \sigma(s)) ds.$$

If

$$b(s) + \beta_s \sigma(s) = 0, \quad \forall s \in [t, T], \quad (14)$$

it follows that the process $(X_s Y_s + \int_t^s (u_r X_r + c_r Y_r) dr)_{s \in [t, T]}$ is a supermartingale (as a positive local martingale), and in particular

$$\mathbb{E} \left[X_T Y_T + \int_t^T (u_s X_s + c_s Y_s) ds \right] \leq xy. \quad (15)$$

Now, by definition of \widetilde{U}_1^* and \widetilde{U}_2 and by (15), if $Y_s > 0$ almost surely for each $s \in [t, T]$, then

$$\begin{aligned} & \mathbb{E} \left[\int_t^T U_1(s, c_s, X_s) ds + U_2(X_T) \right] \\ & \leq \mathbb{E} \left[\int_t^T (\widetilde{U}_1^*(s, Y_s, u_s) + c_s Y_s + u_s X_s) ds + \widetilde{U}_2(Y_T) + X_T Y_T \right] \\ & \leq \mathbb{E} \left[\int_t^T \widetilde{U}_1^*(s, Y_s, u_s) ds + \widetilde{U}_2(Y_T) \right] + xy. \end{aligned} \quad (16)$$

Since $(c, \pi) \in \mathcal{A}(t, x)$ is arbitrary, taking the supremum over $(c, \pi) \in \mathcal{A}(t, x)$ on the left handside in (72), we get for every $u \in \mathcal{U}(t, y)$

$$V(t, x) \leq \mathbb{E} \left[\int_t^T \widetilde{U}_1^*(s, Y_s, u_s) ds + \widetilde{U}_2(Y_T) \right] + xy. \quad (17)$$

Therefore, when (14) holds, the right handside of (17) is an upper bound for the primal value function. On the other hand we can take the infimum over $u \in \mathcal{U}_\beta(t, y)$ in the right handside of (17). Taking into account that (17) has been derived under (14), this leads to consider the control problem

$$(\mathbf{D}) \quad \inf_{u \in \mathcal{U}(t, y)} \widetilde{J}(t, y; u),$$

where $\mathcal{U}(t, y)$ is the set defined in (13) when β is given by (14),

$$\widetilde{J}(t, y; u) = \mathbb{E} \left[\int_t^T \widetilde{U}_1^*(s, Y_s^{t, y, u}, u_s) ds + \widetilde{U}_2(Y_T^{t, y, u}) \right], \quad (18)$$

and $Y^{t, y, u}$ is the solution to (12) when β is given by (14), i.e. the solution to

$$\begin{cases} dY_s = -u_s ds - \frac{b(s)}{\sigma(s)} Y_s dW_s, \\ Y_t = y. \end{cases} \quad (19)$$

We denote by W the value function associated to this problem - that we call *dual value function* - i.e.

$$W(t, y) := \inf_{u \in \mathcal{U}(t, y)} \widetilde{J}(t, y; u), \quad (t, y) \in [0, T] \times (0, +\infty). \quad (20)$$

Taking the infimum over $u \in \mathcal{U}(t, y)$ in the right handside of (17) we get the inequality

$$V(t, x) \leq W(t, y) + xy, \quad \forall (t, y) \in [0, T] \times (0, +\infty). \quad (21)$$

Defining the Legendre transform of the primal value function

$$\widetilde{V}(t, y) := \sup_{x \geq 0} \{V(t, x) - xy\}, \quad (t, y) \in [0, T] \times (0, +\infty),$$

from (21) we get

$$\widetilde{V} \leq W, \quad \text{on } [0, T] \times (0, +\infty). \quad (22)$$

What one can expect is the equality

$$\widetilde{V} = W, \quad \text{on } [0, T] \times (0, +\infty). \quad (23)$$

We will prove (23) as corollary of our next results.

By standard stochastic control arguments we may associate to W an HJB equation that we call dual HJB equation. It is the semilinear equation

$$-w_t(t, y) - \frac{b^2(t)}{2\sigma^2(t)} y^2 w_{yy}(t, y) - \inf_{u \geq 0} \widetilde{H}_{cv}(t, y, w_y(t, y)) = 0, \quad (24)$$

where

$$\widetilde{H}_{cv}(t, y, q) := \widetilde{U}_1^*(t, y, u) - uq,$$

with terminal condition $W(T, \cdot) = \widetilde{U}_2$. Since $U_1^*(t, y, \cdot)$ is concave, the inf-Legendre transform of $\widetilde{U}_1^*(t, y, \cdot)$ on $(-\infty, 0)$, i.e.

$$\inf_{u \geq 0} \{\widetilde{U}_1^*(t, y, \cdot) - uq\}, \quad q < 0,$$

is equal to $U_1^*(t, y, \cdot)$. So, where $w_y < 0$ - it will be the case of our solution - (24) can be rewritten as

$$-w_t(t, y) - \frac{b^2(t)}{2\sigma^2(t)} y^2 w_{yy}(t, y) - U_1^*(t, y, -w_y(t, y)) = 0. \quad (25)$$

Remark 3.1 *Due to the presence of current cost in the state (i.e. the dependence of U_1 on x), we have a (real) dependence of \widetilde{U}_1^* on u_s in the functional (18) defining the dual problem. Since this dependence is monotone (nonincreasing) and since \widetilde{U}_1^* is also nonincreasing on Y_s and u_s appears with the negative sign in (19), this creates a real trade-off between the functional (18) and the state equation (19), giving rise to a real (nontrivial) control problem. At the level of the dual HJB equation (24) above, this can be appreciated by the presence of a nonlinearity in the first order term. When, as in [2, 24], the function U_1 does not depend on x ,⁴ the dependence of this term on w_y disappears and the dual HJB equation is linear. While in [2] the linearity of the dual equation allows to deal with analytical solutions expressed through the heat kernel, a different and more theoretical approach is needed here. We are not aware of papers where the dual problem is investigated when also utility on the current wealth is considered; nevertheless, we stress that utility on the current wealth arises in concrete problems, as the ones described in Section 6.*

4 The dual value function as classical solution of the dual HJB equation

In this section we show that W is a classical solution to the HJB equation (25). To do that first we show that it is a viscosity solution to (25) and then we show its regularity.

4.1 W as viscosity solution of the dual HJB equation

Before to proceed, we need to investigate some properties of \widetilde{U}_1^* , \widetilde{U}_2 and derive qualitative properties for W .

Proposition 4.1 *We have the following properties of the functions \widetilde{U}_1^* and \widetilde{U}_2 .*

1. $\widetilde{U}_1^* : \mathbb{R}_+ \times (0, +\infty) \times (0, +\infty) \rightarrow \mathbb{R}$ is nonnegative, convex in (y, u) and nonincreasing in y and u .

⁴ Actually in [2] the function U_1 expressing the current utility is not even considered. However, as outlined in [2], considering a current utility depending *only* on consumption would not complicate the mathematical problem.

2. $\widetilde{U}_2 : (0, +\infty) \rightarrow \mathbb{R}$ is nonnegative, convex and nonincreasing.

3. We have the following growth estimate: there exists $\tilde{K} > 0$ such that

$$\widetilde{U}_1^*(t, y, u) + \widetilde{U}_2(y) \leq \tilde{K}(1 + y^{-\frac{p}{1-p}} + u^{-\frac{p}{1-p}}), \quad t \in [0, T], \quad u > 0, \quad y > 0. \quad (26)$$

4. We have

$$(i) \quad \lim_{y \wedge u \rightarrow +\infty} \widetilde{U}_1^*(t, y, u) = 0; \quad (ii) \quad \lim_{y \rightarrow +\infty} \widetilde{U}_2(y) = 0. \quad (27)$$

5. According to (a) or (b) of Assumption 2.2(iv), we have respectively either

$$(a) \quad \exists \varepsilon > 0 \text{ such that } \lim_{y \rightarrow 0^+} \widetilde{U}_1^*(t, y, u) = +\infty \text{ uniformly in } (t, u) \in [T - \varepsilon, T] \times \mathbb{R}_+,$$

or

$$(b) \quad \lim_{y \rightarrow 0^+} \widetilde{U}_2(y) = +\infty,$$

or both.

6. $U_1^* \in C_{loc}^{\delta/2, k+\delta}([0, T] \times (0, +\infty) \times (0, +\infty); \mathbb{R})$.

Proof. 1-2-3 follow straightly by using the properties of Legendre transforms and Assumption 2.2(i, ii, iii).

4. For fixed $t > 0$, let for $y > 0, u > 0$,

$$\Lambda_{y,u} = \left\{ (x, c) \in \mathbb{R}_+^2 \mid \frac{\partial}{\partial x} U_1(t, x, c) \geq y, \quad \frac{\partial}{\partial c} U_1(t, x, c) \geq u \right\} \cup \{(0, 0)\}.$$

Using Assumption 2.2(i), it is not difficult to see that the maximizer in the definition of $\widetilde{U}_1^*(t, y, u)$ belongs to $\Lambda_{y,u}$ and that $\Lambda_{y,u}$ shrinks to $\{(0, 0)\}$ as $y \rightarrow +\infty$ and $u \rightarrow +\infty$; so

$$\limsup_{y \wedge u \rightarrow +\infty} \widetilde{U}_1^*(t, y, u) \leq \limsup_{y \wedge u \rightarrow +\infty} \sup_{(x,c) \in \Lambda_{y,u}} U_1(t, x, c) = U_1(t, 0, 0) = 0.$$

The limit for \widetilde{U}_2 follows with a similar argument.

5. If we are in the case of Assumption 2.2(iv)(a), then, due to monotonicity with respect to u of \widetilde{U}_1^* , the statement (a) is equivalent to

$$\exists \varepsilon > 0 \text{ such that } \lim_{y \rightarrow 0} \lim_{u \rightarrow +\infty} \widetilde{U}_1^*(t, y, u) = +\infty, \text{ uniformly w.r.t. } t \in [T - \varepsilon, T]. \quad (28)$$

Now, by (4), using the same argument of point 4 above, but with respect to u only, we get

$$\lim_{u \rightarrow +\infty} \widetilde{U}_1^*(t, y, u) = \sup_{c \geq 0} \{U_1(t, c, 0) - cy\}, \text{ uniformly w.r.t. } t \in [T - \varepsilon, T]. \quad (29)$$

Since taking $c = 1/y$ we get

$$\sup_{c \geq 0} \{U_1(t, c, 0) - cy\} \geq U_1(t, 1/y, 0) - 1, \quad (30)$$

the claim (28) follows combining (29)-(30) and using Assumption 2.2(iv)(a).

In the case of Assumption 2.2(iv)(b) the claim (b) can be obtained as above (but more easily) by using the definition (11).

6. If Assumption 2.2(i)(b) holds, the claim is immediate as

$$U_1^* = U_1 \in C_{loc}^{\delta/2, k+\delta}([0, T] \times (0, +\infty) \times (0, +\infty); \mathbb{R}).$$

Let us prove the claim in the case when Assumption (2.2)(i)(a) holds true. Under our assumptions, the map $c \mapsto \frac{\partial}{\partial c} U_1(t, \cdot, x)$ is a bijection from $(0, +\infty)$ to $(0, +\infty)$ for each $(t, x) \in [0, T] \times (0, +\infty)$, and the supremum in the definition of U_1^* is attained at the unique $c^*(t, y, x)$ satisfying

$$\frac{\partial}{\partial c} U_1(t, c^*(t, y, x), x) = y. \quad (31)$$

Since $\frac{\partial}{\partial c^2} U_1 < 0$, it follows from the implicit function theorem that c^* has the same regularity properties as $\frac{\partial}{\partial c} U_1$, i.e. it is $C_{loc}^{\delta/2, k-1+\delta}([0, T] \times (0, +\infty) \times (0, +\infty); \mathbb{R})$. Writing

$$U_1^*(t, y, x) = U_1(t, c^*(t, y, x), x) - c^*(t, y, x)y$$

and using (31), we obtain

$$\begin{aligned} \frac{\partial}{\partial y} U_1^*(t, y, x) &= c^*(t, y, x), \\ \frac{\partial}{\partial x} U_1^*(t, y, x) &= \frac{\partial}{\partial x} U_1(t, c^*(t, y, x), x). \end{aligned}$$

Both of these functions lie in $C_{loc}^{\delta/2, k-1+\delta}([0, T] \times (0, +\infty) \times (0, +\infty); \mathbb{R})$, which proves the claim. \square

Proposition 4.2 *W is finite, strictly positive on D_T , convex and strictly decreasing in y . Moreover, we have the growth condition, for some $K_W > 0$,*

$$W(t, y) \leq K_W(1 + y^{-\frac{p}{1-p}}), \quad \forall (t, y) \in [0, T] \times (0, +\infty), \quad (32)$$

and terminal and boundary conditions

$$\begin{cases} (i) & W(T, y) = \widetilde{U}_2(y), & \forall y \in (0, +\infty); \\ (ii) & \lim_{y \rightarrow 0^+} W(t, y) = +\infty, & \forall t \in [0, T]; \\ (iii) & \lim_{y \rightarrow +\infty} W(t, y) = 0, & \forall t \in [0, T]. \end{cases} \quad (33)$$

Sketch of proof. The arguments are quite standard and we only sketch the proof of the claims which are straightforward.

Taking the feedback control $u_s = Y_s$ in the state equation (19) and using (26), we obtain that W is finite and satisfies the growth condition (32). The strict positivity in D_T is more tricky and we give a complete proof, which follows from Proposition 4.1(5). Indeed, let $(t, y) \in D_T$. Since $Y_T^{t, y, u} \leq Y_T^{t, y, 0}$ for each $u \in \mathcal{U}(t, y)$, we get

$$\widetilde{J}(t, y; u) \geq \mathbb{E} \left[\int_t^T \widetilde{U}_1^*(s, Y_s^{t, y, 0}, u_s) ds + \widetilde{U}_2(Y_T^{t, y, 0}) \right], \quad \forall u \in \mathcal{U}(t, y). \quad (34)$$

Since $Y^{t,y,0}$ is a Geometric Brownian Motion, setting

$$A_{\varepsilon,y_0}^{t,y} := \left\{ \sup_{s \in [t \vee (T-\varepsilon), T]} Y_s^{t,y,0} < y_0 \right\},$$

we have

$$p_{\varepsilon,y_0}^{t,y} := \mathbb{P}(A_{\varepsilon,y_0}^{t,y}) > 0, \quad \forall \varepsilon > 0, \forall y_0 > 0. \quad (35)$$

Now, set for all $(s, y_0) \in [0, T] \times (0, +\infty)$

$$g(s, y_0) := \lim_{u \rightarrow +\infty} \widetilde{U}_1^*(s, y_0, u). \quad (36)$$

Using (34), (35) and (36), we get

$$\widetilde{J}(t, y; u) \geq p_{\varepsilon,y_0}^{t,y} \left[\int_{t \vee (T-\varepsilon)}^T g(s, y_0) ds + \widetilde{U}_2(y_0) \right], \quad \forall u \in \mathcal{U}(t, y). \quad (37)$$

Now, if Assumption 2.2(iv)(a) holds, take ε above as the one in appearing in the same assumption. By Proposition 4.1(5)(a), we can choose $y_0 > 0$ such that $g(s, y_0) \geq \delta$ for all $s \in [t, T]$ for a suitable $\delta > 0$. Since (37) is uniform in $u \in \mathcal{U}(t, y)$, we get the claim in this case. If we assume that Assumption 2.2(iv)(b) holds, then from it, (37) and Proposition 4.1(5)(b) still follows the claim.

Convexity comes from convexity of \widetilde{U}_1^* and \widetilde{U}_2 , and from linearity of the state equation by standards arguments. Also monotonicity is consequence of standard arguments due to monotonicity of \widetilde{U}_1^* and \widetilde{U}_2 .

The terminal condition (33)(i) comes from the definition of W immediately.

The boundary condition (33)(ii) can be obtained arguing as in the proof of strict positivity of W . Indeed, we can consider (37) with $y_0 = y$. Then, since $Y^{t,y,0} = yY^{t,1,0}$ we get that

$$p_{\varepsilon,y}^{t,y} = p_{\varepsilon,1}^{t,1} > 0, \quad \forall y \in (0, 1). \quad (38)$$

Therefore, (37) becomes in this case

$$\widetilde{J}(t, y; u) \geq p_{\varepsilon,1}^{t,1} \left[\int_{t \vee (T-\varepsilon)}^T g(s, y) ds + \widetilde{U}_2(y) \right], \quad \forall u \in \mathcal{U}(t, y). \quad (39)$$

from which we get

$$W(t, y) \geq p_{\varepsilon,1}^{t,1} \left[\int_{t \vee (T-\varepsilon)}^T g(s, y) ds + \widetilde{U}_2(y) \right]. \quad (40)$$

Taking the limit for $y \rightarrow 0^+$ and using Proposition 4.1(5), we get (33)(ii).

Let us show now the boundary condition (33)(iii). Let $(t, y) \in [0, T] \times (0, +\infty)$ and take the feedback control $u_s = Y_s$ in (19) and consider the associated state trajectory $Y_s^{t,y,u}$. Then

$$W(t, y) \leq \widetilde{J}(t, y; u) = \mathbb{E} \left[\int_t^T \widetilde{U}_1^*(s, Y_s^{t,y,u}, Y_s^{t,y,u}) ds + \widetilde{U}_2(Y_T^{t,y,u}) \right]. \quad (41)$$

Since

$$Y_s^{t,y,u} = y \cdot \exp \left(- \int_t^s \left(1 + \frac{b^2(\xi)}{2\sigma^2(\xi)} \right) d\xi - \int_t^s \frac{b(\xi)}{\sigma(\xi)} dW_\xi \right),$$

we have

$$Y_s^{t,y,u} \rightarrow +\infty, \quad \forall s \in [t, T], \text{ a.s.} \quad (42)$$

Hence, using (27) and (42) we get

$$\widetilde{U}_1^*(s, Y_s^{t,y,u}, Y_s^{t,y,u}) \rightarrow 0, \quad \forall s \in [t, T], \text{ a.s., and } \quad \widetilde{U}_2(Y_T^{t,y,u}) \rightarrow 0, \text{ a.s.} \quad (43)$$

On the other hand, thanks to (26), we have

$$\widetilde{U}_1^*(s, Y_s^{t,y,u}, Y_s^{t,y,u}) \leq \tilde{K} \left(1 + 2(Y_s^{t,y,u})^{-\frac{p}{1-p}} \right), \quad \widetilde{U}_2(Y_T^{t,y,u}) \leq \tilde{K} (1 + (Y_T^{t,y,u})^{-\frac{p}{1-p}}).$$

Since the above right hand sides are integrable uniformly in $y \geq 1$, using (41) and (43) we get the claim by Vitali's Theorem.

Finally, strict monotonicity follows from convexity, monotonicity, strict positivity and (33)(iii). \square

Proposition 4.3 *W is continuous on $[0, T] \times (0, +\infty)$. Moreover $W(\cdot, y)$ is nondecreasing for all $y \in (0, +\infty)$.*

Proof. First of all, by convexity, W is continuous in the space variable y for each fixed $t \in [0, T]$.

Let us show continuity in time. For that, we need to exploit the following Dynamic Programming Principle:⁵ for each t, t' such that $0 \leq t \leq t' \leq T$ and each $y \in (0, +\infty)$,

$$W(t, y) = \inf_{u \in \mathcal{U}(t, y)} \mathbb{E} \left[\int_t^{t'} \widetilde{U}_1^*(s, Y_s^{t,y,u}, u_s) ds + W(t', Y_{t'}^{t,y,u}) \right]. \quad (44)$$

Now we show that W is nonincreasing in time. Indeed, let $(t, y) \in [0, T] \times (0, +\infty)$, let $u \in \mathcal{U}(t, y)$ and let $t' \in [t, T]$. Since $\widetilde{U}_1^* \geq 0$, from (44) we have

$$W(t, y) \geq \inf_{u \in \mathcal{U}(t, y)} \mathbb{E} \left[W(t', Y_{t'}^{t,y,u}) \right]. \quad (45)$$

By monotonicity of W in y and since $Y_{t'}^{t,y,u} \leq Y_{t'}^{t,y,0}$ for all $u \in \mathcal{U}(t, y)$, we get

$$\inf_{u \in \mathcal{U}(t, y)} \mathbb{E} \left[W(t', Y_{t'}^{t,y,u}) \right] \geq \mathbb{E} \left[W(t', Y_{t'}^{t,y,0}) \right]. \quad (46)$$

Combining (45) and (46), and using Jensen's inequality, we finally get

$$W(t, y) \geq W(t', y),$$

⁵Appealing to the Dynamic Programming Principle may seem somehow unfair, as usually it is problematic to prove it if one has not proved before the continuity of the value function (and we are just proving the continuity invoking it). However, we observe that in this case (where the time t' is deterministic) the proof of the Dynamic Programming Principle (see, e.g., [25, Ch. 4]), only uses the continuity in the space variable y .

proving the monotonicity claim.

From this monotonicity it follows that the functions provided by the left and right limits of W in t , i.e.

$$W_+(t, \cdot) := \lim_{h \downarrow 0} W(t+h, \cdot), \quad W_-(t, \cdot) := \lim_{h \downarrow 0} W(t-h, \cdot),$$

are well-defined in $[0, T)$ and $(0, T]$ respectively, and

$$W_- \geq W \geq W_+ \quad (47)$$

(where the functions are defined). We note that W_+, W_- are also convex in y for fixed t , so they are continuous in y for fixed t as well. If we show the inequalities

$$W_- \leq W \leq W_+ \quad (48)$$

(where the functions are defined) combining with (47) the proof of continuity in time will be complete.

Let us first show the left inequality in (48). For any $s \in [0, T]$, define $\hat{Y}^{s,y}$ as the process corresponding to the feedback control $\hat{u} = \hat{Y}$ starting from (s, y) . Then, for each $r \geq s$,

$$\hat{Y}_r^{s,y} = y \exp \left(\int_s^r \left(-1 + \frac{1}{2} \frac{b(\xi)^2}{\sigma(\xi)^2} \right) d\xi - \int_s^r \frac{b(\xi)}{\sigma(\xi)} dW_\xi \right).$$

Note that, since $\frac{b(\cdot)}{\sigma(\cdot)}$ is bounded, we have the following estimates :

$$\mathbb{E} \left[\left| \hat{Y}_r^{s,y} - y \right| \right] \leq \omega(|s-r|), \quad \text{with } \omega \text{ continuous and } \omega(0^+) = 0, \quad (49)$$

$$\sup_{0 \leq r \leq s \leq T} \mathbb{E} \left[\left| \hat{Y}_r^{s,y} \right|^q \right] < +\infty, \quad \forall q \in \mathbb{R}. \quad (50)$$

Let $t \in [0, T]$ and take a sequence $t_n \uparrow t$. By (44) and (26),

$$\begin{aligned} W(t_n, y) &\leq \mathbb{E} \left[\int_{t_n}^t \widetilde{U}_1^*(s, \hat{Y}_s^{t_n,y}, \hat{Y}_s^{t_n,y}) ds + W(t, \hat{Y}_t^{t_n,y}) \right] \\ &\leq \mathbb{E} \left[\int_{t_n}^t 2\tilde{K} (1 + (\hat{Y}_s^{t_n,y})^{-\frac{p}{1-p}}) ds + W(t, \hat{Y}_t^{t_n,y}) \right]. \end{aligned} \quad (51)$$

By (50) the expectation of the integral in (51) goes to 0. On the other hand, from (49), passing to a subsequence if necessary (we have monotonicity in t , so we can do that without loss of generality), we see that $\hat{Y}_t^{t_n,y} \rightarrow y$ almost surely. Hence, using (50) and the growth condition (32) on W , by dominated convergence we get

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[W(t, \hat{Y}_t^{t_n,y}) \right] = W(t, y).$$

So, we finally obtain $W_-(t, y) \leq W(t, y)$.

Now let us turn to the proof of the right inequality in (48). Let $t \in [0, T]$ and take a sequence $t_n \downarrow t$. Again, using (44) we have that

$$W(t, y) \leq \mathbb{E} \left[\int_t^{t_n} \widetilde{U}_1^*(s, \hat{Y}_s^{t,y}, \hat{Y}_s^{t,y}) ds + W(t_n, \hat{Y}_{t_n}^{t,y}) \right].$$

The proof is now the same once we show that $W(t_n, \hat{Y}_{t_n}^{t,y}) \rightarrow W(t^+, y)$ almost surely. We observe that $W(t_n, \cdot) \searrow W_+(t, \cdot)$ pointwise by definition. Since all these functions are continuous, by Dini's Theorem we get $W(t_n, \cdot) \searrow W_+(t, \cdot)$ locally uniformly. Therefore $t_n \downarrow t$, $y_n \rightarrow y$ implies $W(t_n, y_n) \rightarrow W_+(t, y)$. Since, by passing to a subsequence if necessary (again we may do that without loss of generality because of monotonicity in t) we can assume $Y_{t_n}^{t,y} \rightarrow y$ almost surely, it follows that $W(t_n, \hat{Y}_{t_n}^{t,y}) \rightarrow W_+(t, y)$ almost surely. And again by dominated convergence this implies $W(t, y) \leq W_+(t, y)$. This completes the proof of continuity in time.

Now it just remains to notice that again by Dini's Theorem the continuity of W in t is locally uniform in y , which combined to the fact that W is continuous in y for fixed t , implies joint continuity of W in (t, y) . \square

Now we may state the viscosity property of W .

Proposition 4.4 *W is a continuous viscosity solution to (25) in D_T .*

Proof. Due to continuity of W , this is quite standard. We omit the proof for brevity and refer to classical references, such as [14, 25]. \square

4.2 Regularity of W

In this section we prove a regularity result for the dual value function W .

Theorem 4.5

1. $W \in C_{loc}^{1+\delta/2, k+2+\delta}(D_T; \mathbb{R})$.
2. $W_y(t, \cdot) < 0$, $W_y(t, 0^+) = -\infty$ and $W_y(t, +\infty) = 0$, for every $t \geq 0$.
3. $W_{yy} > 0$ over D_T .

Proof. 1. Take any $(t_0, y_0) \in D_T$ and consider, for suitable $\varepsilon > 0$, the square

$$D_\varepsilon(t_0, y_0) := [t_0, t_0 + \varepsilon) \times (y_0 - \varepsilon, y_0 + \varepsilon) \subset D_T.$$

First of all, note that, due to convexity, the right and left space derivatives of W exist. Denoting them by $W_y(t, y^+)$ and $W_y(t, y^-)$ respectively, again by convexity we have $W_y(t, y^+) \geq W_y(t, y^-)$. Moreover, there exist $M_\varepsilon, m_\varepsilon > 0$ such that

$$M_\varepsilon \geq \sup_{(t,y) \in D_\varepsilon(t_0, y_0)} -W_y(t, y^-) \geq \inf_{(t,y) \in D_\varepsilon(t_0, y_0)} -W_y(t, y^+) \geq m_\varepsilon. \quad (52)$$

Indeed, by convexity $-W_y(t, y^+) \geq \frac{1}{y}(W(t, y) - W(t, 2y))$, and, since W is continuous and strictly decreasing in y for each t , the infimum above must be strictly positive. In the same way, $-W_y(t, y^-) \leq \frac{2}{y}(W(t, y) - W(t, y/2))$ and the supremum is finite.

By Proposition 4.4, the dual value function W is a viscosity solution of the dual HJB equation (25) in $D_\varepsilon(t_0, y_0)$ with Dirichlet boundary condition

$$w = W, \quad \text{on } \mathcal{P}(D_\varepsilon(t_0, y_0)), \quad (53)$$

where $\mathcal{P}(D_\varepsilon(t_0, y_0))$ is the parabolic boundary of $D_\varepsilon(t_0, y_0)$ defined as

$$\mathcal{P}(D_\varepsilon(t_0, y_0)) := \{t_0 + \varepsilon\} \times [y_0 - \varepsilon, y_0 + \varepsilon] \cup [t_0, t_0 + \varepsilon] \times \{y_0 - \varepsilon, y_0 + \varepsilon\}.$$

Consider the function F defined on $D_\varepsilon(t_0, y_0) \times \mathbb{R}$ by

$$F(t, y, q) := U_1^*(t, y, -[(m_\varepsilon \vee q) \wedge M_\varepsilon]).$$

By Proposition 4.1(6), F is Hölder continuous in $D_\varepsilon(t_0, y_0) \times \mathbb{R}$. By (52), we have that W is actually a viscosity solution in $D_\varepsilon(t_0, y_0)$ to the equation

$$-w_t(t, y) - \frac{b^2(t)}{2\sigma^2(t)} y^2 w_{yy}(t, y) - F(t, y, w_y(t, y)) = 0. \quad (54)$$

Since W is continuous on $\mathcal{P}(D_\varepsilon(t_0, y_0))$, then we have uniqueness of viscosity solutions to (54) with boundary condition (53) (see, e.g., [14, Cor. 8, Ch. V]). On the other hand, due to Assumption 2.1 and to Hölder continuity of F , the PDE (54) is semilinear uniformly parabolic on $D_\varepsilon(t_0, y_0)$ with Hölder continuous coefficients, so by Theorem 12.22 of [22] - with the assumptions of Theorem 12.16 of the same book - it admits a solution fulfilling the boundary condition (53) in the space $C^{1,2}(D_\varepsilon(t_0, y_0); \mathbb{R})$. This (classical) solution is also a viscosity solution, thus, due to uniqueness of viscosity solutions, it coincides with W . Hence, we conclude that $W \in C^{1,2}(D_\varepsilon(t_0, y_0); \mathbb{R})$, therefore, by arbitrariness of (t_0, y_0) , that $W \in C^{1,2}(D_T; \mathbb{R})$.

Given that, we know that $-W_y$ is strictly positive and locally Lipschitz continuous in D_T . Moreover, by Proposition 4.1(6), $U_1^* \in C_{loc}^{\delta/2, k+\delta}([0, T) \times (0, +\infty) \times (0, +\infty); \mathbb{R})$. Therefore, the claim follows from a simple induction, using regularity results for linear equations of the form $-u_t - Lu = f$ (see, e.g., Theorem 8.12.1, p. 131, in [21]).

2. The first claim follows (52). The other ones follow from convexity and from (33)(ii) and (33)(iii), respectively.

3. As in [2] we use a maximum principle argument. Differentiating twice (25), we get

$$\begin{aligned} - (W_{yy})_t - \frac{b^2(s)}{2\sigma^2(s)} [2W_{yy} + 4y(W_{yy})_y + y^2(W_{yy})_{yy}] \\ - (U_1^*)_{yy}(t, y, -W_y) + W_{yyy} \cdot (U_1^*)_x(t, y, -W_y) \\ + 2W_{yy} \cdot (U_1^*)_{xy}(t, y, -W_y) - W_{yy}^2 \cdot (U_1^*)_{xx}(t, y, -W_y) = 0. \end{aligned}$$

Noting that U_1^* is convex in y , we see that W_{yy} is a nonnegative supersolution to the linear parabolic PDE

$$\begin{aligned} -u_t - \frac{b^2(s)}{2\sigma^2(s)} [2u + 4yu_y + y^2u_{yy}] + (U_1^*)_x(t, y, -W_y)u_y \\ + [2(U_1^*)_{xy}(t, y, -W_y) - W_{yy} \cdot (U_1^*)_{xx}(t, y, -W_y)]u = 0. \end{aligned}$$

Hence, by a strong maximum principle (see e.g. [15, Th. 3, Ch. II]), if $W_{yy}(t_0, y_0) = 0$ for some $(t_0, y_0) \in D_T$, it must be $W_{yy} \equiv 0$ on $(t_0, T) \times (0, +\infty)$, which is clearly in contradiction, e.g., with (33)(ii). \square

From Proposition 4.4 and Theorem 4.5 we get the following

Corollary 4.6 *W is a classical solution to (25) in D_T .*

5 Back to the primal control problem: verification and optimal controls

Let $t \in [0, T]$ and let \widetilde{W} be the inf-Legendre transform of $W(t, \cdot)$, i.e.

$$\widetilde{W}(t, x) := \inf_{y > 0} \{W(t, y) + xy\}, \quad (t, x) \in \overline{D_T}. \quad (55)$$

Due to its definition and to the positivity of W (see Proposition 4.2), the function \widetilde{W} is finite and nonnegative on $\overline{D_T}$. Moreover, it is concave and nondecreasing in x for each $t \in [0, T]$ and, due to Theorem 4.5, it can be written, for $(t, x) \in D_T$, as

$$\widetilde{W}(t, x) = W(t, [W_y(t, \cdot)]^{-1}(-x)) + x [W_y(t, \cdot)]^{-1}(-x). \quad (56)$$

We are going to prove that

$$\widetilde{W} = V, \quad \text{on } \overline{D_T} \quad (57)$$

(we notice that (57) implies, as corollary, (23), i.e. $\widetilde{V} = W$) and that V is the unique classical solution of the primal HJB equation (8) in the following class:

$$\begin{aligned} \mathcal{C} = \Big\{ & v \in C(\overline{D_T}; \mathbb{R}) \cap C_{loc}^{1+\delta/2, k+2+\delta}(D_T; \mathbb{R}) \text{ such that } v_x > 0, v_{xx} < 0 \text{ in } D_T, \\ & \text{and } v \text{ fulfills the boundary and growth conditions (58) below} \Big\} \end{aligned}$$

where

$$\begin{cases} (i) & v(t, 0) = 0, \quad \forall t \in [0, T], \\ (ii) & v(T, x) = U_2(x), \quad \forall x \geq 0, \\ (iii) & \exists K_0 \text{ such that } 0 \leq v(t, x) \leq K_0(1 + x^p), \quad \forall (t, x) \in [0, T] \times [0, +\infty). \end{cases} \quad (58)$$

We note that if $v \in \mathcal{C}$, due to (9), we have

$$H(t, x, v_x(t, x), v_{xx}(t, x)) = U_1^*(t, v_x(t, x), x) - \frac{b^2(t)}{2\sigma^2(t)} \frac{v_x(t, x)^2}{v_{xx}(t, x)}. \quad (59)$$

We proceed as follows:

1. We show that $\widetilde{W} \in \mathcal{C}$ and that it is a classical solution of the primal HJB equation (8) (Proposition 5.1).
2. We show that a verification theorem holds for **(P)** for every classical solution $v \in \mathcal{C}$ of the primal HJB equation (Theorem 5.2).
3. We show that for every classical solution $v \in \mathcal{C}$ of the primal HJB equation the associated closed loop equation admits a solution and that this implies $v = V$ (Proposition 5.3 and Corollary 5.4).

Clearly, these three points yield the equality $\widetilde{W} = V$ and the announced uniqueness.

5.1 \widetilde{W} is a classical solution of the primal HJB equation

Proposition 5.1 $\widetilde{W} \in \mathcal{C}$ and solves the primal HJB equation (8) in classical sense in D_T . Moreover it satisfies the Inada conditions in x :

$$\widetilde{W}_x(t, 0^+) = +\infty, \quad \widetilde{W}_x(t, +\infty) = 0, \quad \forall t \in [0, T].$$

Proof. *Growth and boundary conditons.* The growth condition (58)(iii) follows from (55) and (32). The boundary condition (58)(i) follows from (55) and (33)(iii). The boundary condition (58)(ii) follows from (55), (33)(i) and the fact that the inf-Legendre transform of \widetilde{U}_2 is U_2 .

Continuity in $\overline{D_T}$. The fact that \widetilde{W} is continuous in D_T follows from (56) and Theorem 4.5. Now we show the continuity at the boundary $[0, T) \times \mathbb{R}$.

Continuity of $\widetilde{W}(t, \cdot)$ at 0^+ for each $t \in [0, T)$ follows from (55): it yields

$$\widetilde{W}(t, x) \leq W(t, \varepsilon/x) + \varepsilon, \quad \forall x \geq 0, \quad \forall \varepsilon > 0,$$

hence, taking into account also that W is nonnegative and (33)(iii),

$$0 \leq \limsup_{x \downarrow 0} \widetilde{W}(t, x) \leq \varepsilon, \quad \forall \varepsilon > 0,$$

and, since ε is arbitrary and taking into account (58)(i), we may conclude that

$$\lim_{x \downarrow 0} \widetilde{W}(t, x) = 0 = \widetilde{W}(t, 0).$$

Moreover, by monotonicity of $\widetilde{W}(t, \cdot)$ for all $t \in [0, T)$ the convergence above is locally uniform in $t \in [0, T)$ due to Dini's Theorem, so, combining with the obvious continuity of $\widetilde{W}(\cdot, 0)$, we get the continuity of \widetilde{W} at the boundary $[0, T) \times \mathbb{R}$ in the couple (t, x) .

Next we show the continuity at the boundary $\{T\} \times \mathbb{R}$. First let us show the continuity of $W(\cdot, x)$ at T^- for fixed $x \in \mathbb{R}$. Since $\widetilde{W}(t, 0) = 0$ for every $t \in [0, T]$, the claim is obvious for $x = 0$, so we now assume $x > 0$. Clearly, for any $y > 0$,

$$\limsup_{t \uparrow T} \widetilde{W}(t, x) \leq \limsup_{t \uparrow T} \{W(t, y) + xy\} = W(T, y) + xy,$$

by continuity of W . Taking the infimum over y , we obtain the inequality

$$\limsup_{t \uparrow T} \widetilde{W}(t, x) \leq \widetilde{W}(T, x).$$

For the opposite inequality, we notice that, by definition of W , we have for each $y > 0$ and each $t \in [0, T]$

$$W(t, y) \geq \mathbb{E} \left[\widetilde{U}_2(Y_T^{t, y, 0}) \right] \geq \widetilde{U}_2 \left(\mathbb{E} \left[Y_T^{t, y, 0} \right] \right) = \widetilde{U}_2(y),$$

where we have used Jensen's inequality. Since $\widetilde{W}(T, \cdot) = U_2(\cdot)$, we get $W(t, \cdot) \geq W(T, \cdot)$, which in turn yields

$$\liminf_{t \uparrow T} \widetilde{W}(t, x) \geq \widetilde{W}(T, x).$$

Now, taking into account the obvious continuity of $\widetilde{W}(T, \cdot)$ in \mathbb{R}_+ , the continuity of \widetilde{W} at the boundary $\{T\} \times \mathbb{R}_+$ in the couple (t, x) follows again from Dini's Theorem, as $\widetilde{W}(\cdot, x)$ inherits from $W(\cdot, y)$ the monotonicity (Proposition 4.3). This concludes the proof of the continuity of \widetilde{W} on $\overline{D_T}$.

Further regularity in D_T . From (56) and taking into account Theorem 4.5, we get for each $(t, y) \in D_T$

$$\begin{cases} (i) & \widetilde{W}_t(t, x) = W_t(t, [W_y(t, \cdot)]^{-1}(-x)), \\ (ii) & \widetilde{W}_x(t, x) = [W_y(t, \cdot)]^{-1}(-x), \\ (iii) & \widetilde{W}_{xx}(t, x) = -\frac{1}{W_{yy}(t, [W_y(t, \cdot)]^{-1}(-x))}. \end{cases} \quad (60)$$

So, due to Theorem 4.5, we have $\widetilde{W} \in C_{loc}^{1+\delta/2, k+2+\delta}(D_T; \mathbb{R})$ and $\widetilde{W}_x > 0, \widetilde{W}_{xx} < 0$ in D_T . This completes the proof that $\widetilde{W} \in \mathcal{C}$.

\widetilde{W} as solution to the HJB equation. The fact that \widetilde{W} solves the HJB equation (8) in classical sense in D_T follows from Corollary 4.6 by straightforward computations using (59) and (60).

Inada's conditions. Inada's conditions follow from Theorem 4.5(2) and (60)(ii). \square

5.2 Verification theorem

Theorem 5.2 *Let $v \in \mathcal{C}$ be a classical solution to the primal HJB equation (8). Then:*

- (i) $v(t, x) \geq V(t, x)$ for all $(t, x) \in \overline{D_T}$.
- (ii) Let $(t, x) \in \overline{D_T}$, let $(c^*, \pi^*) \in \mathcal{A}(t, x)$ and let $X^* := X^{t, x, c^*, \pi^*}$. If

$$H_{cv}(s, X_s^*, v_x(s, X_s^*), v_{xx}(s, X_s^*); c_s^*, \pi_s^*) = H(s, X_s^*, v_x(s, X_s^*), v_{xx}(s, X_s^*)) \quad (61)$$

\mathbb{P} -almost surely for almost every $s \in [t, T]$, then (c^*, π^*) is an optimal control and $v(t, x) = V(t, x)$.

Proof. (i) Let $(t, x) \in \overline{D_T}$, $(c, \pi) \in \mathcal{A}(t, x)$, and, to simplify the notation, let us write $X_s := X_s^{t, x, c, \pi}$ for all $s \in [t, T]$. Set

$$\tau := \inf \{s \in [t, T] \mid X_s = 0\} \wedge T.$$

We notice that, due to the state constraint, $\mathcal{A}(s, 0) = \{(0, 0)\}$ for all $s \in [t, T]$ and the corresponding state trajectory is identically 0, so

$$\text{if } \tau < T, \text{ then } (c, \pi, X) \equiv (0, 0, 0) \text{ in the random time interval } [\tau, T]. \quad (62)$$

Now we may find a sequence of stopping times $\tau_n \nearrow \tau$ such that $v_x(\cdot, X_\cdot)$ is a martingale in $[t, \tau_n]$. Since $v \in C^{1,2}([t, T] \times (0, +\infty); \mathbb{R})$ and satisfies the HJB equation (8), Itô's formula

yields

$$\begin{aligned}\mathbb{E}[v(\tau_n, X_{\tau_n})] &= v(t, x) + \mathbb{E}\left[\int_t^{\tau_n} (H_{cv} - H)(s, X_s, v_x(s, X_s), v_{xx}(s, X_s); c_s, \pi_s) ds\right] \\ &\quad - \mathbb{E}\left[\int_t^{\tau_n} U_1(s, c_s, X_s) ds\right] \\ &\leq v(t, x) - \mathbb{E}\left[\int_t^{\tau_n} U_1(s, c_s, X_s) ds\right].\end{aligned}$$

This gives us

$$v(t, x) \geq \mathbb{E}\left[v(\tau_n, X_{\tau_n}) + \int_t^{\tau_n} U_1(s, c_s, X_s) ds\right], \quad \forall n \in \mathbb{N}. \quad (63)$$

Letting $n \rightarrow \infty$ in (63), using Fatou's Lemma on the first term of the expectation of the right handside, and monotone convergence on the second one, we get

$$\begin{aligned}v(t, x) &\geq \mathbb{E}\left[v(\tau, X_\tau) + \int_t^\tau U_1(s, c_s, X_s) ds\right] \\ &= \mathbb{E}\left[\mathbf{1}_{\{\tau < T\}} \left(v(\tau, X_\tau) + \int_t^\tau U_1(s, c_s, X_s) ds\right)\right] \\ &\quad + \mathbb{E}\left[\mathbf{1}_{\{\tau = T\}} \left(v(\tau, X_\tau) + \int_t^\tau U_1(s, c_s, X_s) ds\right)\right].\end{aligned} \quad (64)$$

Using (62), the fact that $U_2(0) = 0$ and that $U_1(\cdot, 0, 0) = v(\cdot, 0) = 0$, we get

$$v(t, x) \geq \mathbb{E}\left[U_2(X_T) + \int_t^T U_1(s, c_s, X_s) ds\right]. \quad (65)$$

Since $(c, \pi) \in \mathcal{A}(t, y)$ was arbitrary, this means that $v(t, x) \geq V(t, x)$, and (i) is proved.

(ii) Let $(c^*, \pi^*) \in \mathcal{A}(t, y)$ satisfy (61), and denote $X^* = X^{t, y, c^*, \pi^*}$. In this case we have equality in (63), i.e.

$$v(t, x) = \mathbb{E}\left[v(\tau_n, X_{\tau_n}^*) + \int_t^{\tau_n} U_1(s, c_s^*, X_s^*) ds\right], \quad \forall n \in \mathbb{N}. \quad (66)$$

Now we take the limit for $n \rightarrow \infty$ keeping the equality above. We cannot use Fatou's Lemma as before for the part $v(\tau_n, X_{\tau_n}^*)$, but we need to use a result keeping the equality in the limit. Since $\lim_{n \rightarrow \infty} v(\tau_n, X_{\tau_n}^*) = v(\tau, X_\tau^*)$ almost surely, it suffices to prove uniform integrability of $(v(\tau_n, X_{\tau_n}^*))_{n \geq 0}$. For this purpose, write $Y_s := Y_s^{t, 1, 0}$ for all $s \in [t, T]$. We know from the discussion following (14) that $(X_s^* Y_s + \int_t^s c_u^* Y_u du)_{s \in [t, T]}$ is a supermartingale. Since $c_s^* Y_s \geq 0$, we see that also $(X_s^* Y_s)_{s \in [0, T]}$ is a supermartingale, hence $\mathbb{E}[X_{\tau_n}^* Y_{\tau_n}] \leq x$. Now, taking $q \in (p, 1)$, we get, using (58)(iii),

$$\mathbb{E}\left[v(\tau_n, X_{\tau_n}^*)^{q/p}\right] \leq \mathbb{E}\left[K_0^{q/p} (1 + |X_{\tau_n}^*|^p)^{q/p}\right] \leq K_0^{q/p} 2^{\frac{q}{p}-1} (1 + \mathbb{E}[|X_{\tau_n}^*|^q]).$$

Now, using Hölder's inequality, from the inequality above we get

$$\mathbb{E}\left[v(\tau_n, X_{\tau_n}^*)^{q/p}\right] \leq K_0 (1 + \mathbb{E}[X_{\tau_n}^* Y_{\tau_n}]^q \mathbb{E}[(Y_{\tau_n})^{-\frac{q}{1-q}}]^{1-q}) \leq K_0' (1 + x^q).$$

So the sequence $v(\tau_n, X_{\tau_n}^*)_{n \geq 0}$ is bounded in $L^{q/p}$ with $q/p > 1$. By de La Vallée Poussin's Theorem it is uniformly integrable. Hence taking the limit in (66) we get

$$v(t, x) = \mathbb{E} \left[v(\tau, X_\tau^*) + \int_t^\tau U_1(s, c_s^*, X_s^*) ds \right]. \quad (67)$$

Splitting on the sets $\{\tau < T\}$ and $\{\tau = T\}$ as above, taking into account that $v(T, \cdot) = U_2(\cdot)$ for the part corresponding to set $\{\tau = T\}$, taking into account (62) and that $v(\cdot, 0) = 0 = U_1(\cdot, 0, 0)$ on the set $\{\tau < T\}$, we finally rewrite (67) as

$$v(t, y) = J(t, y; c^*, \pi^*). \quad (68)$$

Combining (68) with the claim (i) we get the claim (ii). \square

From Proposition 5.1 and Theorem 5.2, we see that $\widetilde{W} \geq V$.⁶ What we want to get is indeed the equality, and in order to get it we need to exploit further item (ii) of Theorem 5.2 finding optimal feedback controls.

5.3 Optimal feedback controls

Given $v \in \mathcal{C}$, we may define feedback maps in classical sense associated to the maximization of H_{cv} in the HJB equation (8). They are, for $s \in [0, T]$,

$$C^v(s, x) = \begin{cases} \begin{cases} \left[\frac{\partial}{\partial c} U_1(t, \cdot, x) \right]^{-1} (v_x(t, x)), & \text{if } x > 0, \\ 0, & \text{if } x = 0, \end{cases} & \text{if Assumption 2.2(i)(a) holds,} \\ 0, & \text{if Assumption 2.2(i)(b) holds,} \end{cases} \quad (69)$$

$$\Pi^v(s, x) = \begin{cases} -\frac{b(s)v_x(s, z)}{\sigma(s)v_{xx}(s, z)}, & \text{if } x > 0, \\ 0, & \text{if } x = 0. \end{cases} \quad (70)$$

Their definition for $x > 0$ is indeed given by the maximization of H_{cv} in the HJB equation taking into account the structure of the Hamiltonian (9) for functions in \mathcal{C} , while the definition at $x = 0$ is due to the state constraint, which implies $\mathcal{A}(t, 0) = \{(0, 0)\}$.

The closed loop equation associated to the feedback maps C^v, Π^v is

$$\begin{cases} dX_s = -C^v(s, X_s)ds + b(s)\Pi^v(s, X_s)ds + \sigma(s)\Pi^v(s, X_s)dW_s, \\ X_t = x. \end{cases} \quad (71)$$

Since $v \in \mathcal{C}$, one has local Lipschitz continuity of $\Pi^v(s, \cdot)$ on $(0, +\infty)$ for every $s \in [t, T]$. and local Lipschitz continuity of $C^v(s, \cdot)$ on $(0, +\infty)$ for every $s \in [t, T]$. So we can prove the following.

⁶This inequality may be also proved using (22) and the concavity of V in x which could be proved directly.

Proposition 5.3 *Given $v \in \mathcal{C}$ and $(t, x) \in [0, T] \times \mathbb{R}_+$, there exists a unique solution $X^{t,x;v} \geq 0$ to the closed loop equation (71) in the interval $[t, T]$.*

Proof. *Existence.* If $x = 0$ the claim is clear, just by taking $X^{t,x;v} \equiv 0$. Let $x > 0$. Due to local Lipschitz continuity of $\tilde{C}^v(s, \cdot), \tilde{\Pi}^v(s, \cdot)$, using standard SDE's theory (see, e.g., [20, Ch. 5, Th. 2.9]), we get for each $\varepsilon > 0$ the existence of a unique solution $X^{t,x,\varepsilon;v} \in [\varepsilon, \varepsilon^{-1}]$ in the stochastic interval $[t, \tau_\varepsilon)$, where τ_ε is implicitly defined in terms of the solution itself as

$$\tau_\varepsilon = \inf \{s \in [t, T] \mid X_s^{t,x,\varepsilon;v} \leq \varepsilon \text{ or } X_s^{t,x,\varepsilon;v} \geq \varepsilon^{-1}\},$$

with the convention $\inf \emptyset = T$. Of course, if $\varepsilon < \varepsilon'$, we have $\tau_\varepsilon > \tau_{\varepsilon'}$ and

$$X_s^{t,x,\varepsilon} \equiv X_s^{t,x,\varepsilon'} \text{ on } [t, \tau_{\varepsilon'}), \quad \forall 0 < \varepsilon < \varepsilon'. \quad (72)$$

Set

$$\tau = \lim_{\varepsilon \downarrow 0} \tau_\varepsilon.$$

Then by (72) there exists a unique solution $X^{t,x;v} \geq 0$ to (71) in the interval $[t, \tau)$. We now show that this solution can be extended to the whole interval $[t, T]$. By a Girsanov transformation, there exists a probability \mathbb{Q} equivalent to \mathbb{P} , and a \mathbb{Q} -Brownian motion \tilde{W} , such that (71) may be rewritten as

$$dX_s = -C^v(s, X_s)ds + \sigma(s)\Pi^v(s, X_s)d\tilde{W}_s.$$

By nonnegativity of \tilde{C}^v and $X^{t,x;v}$, the process $X^{t,x;v}$ is a nonnegative \mathbb{Q} -supermartingale on $[t, \tau)$. It can be extended to a \mathbb{Q} -supermartingale (L^1 bounded) on $[t, T]$ setting $X^{t,x;v} \equiv 0$ in $[\tau, T]$. Hence, by Doob's convergence Theorem (the usual proof for deterministic intervals - see e.g. Theorem 6.18 in [18] - can be easily adapted to our stochastic interval $[t, \tau)$), there exists a finite random variable $X_\tau^{t,x;v}$ such that

$$\lim_{s \nearrow \tau} X_s^{t,x;v} = X_\tau^{t,x;v}, \quad \mathbb{Q}\text{-a.s.} \quad (73)$$

Since $\mathbb{Q} \sim \mathbb{P}$, we also have

$$\lim_{s \nearrow \tau} X_s^{t,x;v} = X_\tau^{t,x;v}, \quad \mathbb{P}\text{-a.s.} \quad (74)$$

Immediately (74) yields the desired extension on $\{\tau = T\}$. Let us now consider the set $\{\tau < T\}$. On this set we have $X_{\tau_\varepsilon}^{t,x;v} \in \{\varepsilon, \varepsilon^{-1}\}$, so that by (74) necessarily $X_\tau^{t,x;v} = 0$ almost surely, getting

$$\lim_{s \nearrow \tau} X_s^{t,x;v} = 0 \quad \text{a.s. on } \{\tau < T\}. \quad (75)$$

Therefore, we may now extend $X^{t,x;v}$ to a solution defined over $[t, T]$ on $\{\tau < T\}$ by setting

$$X_s^{t,x;v} \equiv 0, \quad \text{for } s \in [\tau, T].$$

Uniqueness. The solution is clearly unique on the stochastic interval $[t, \tau)$ defined in the existence part. If it reaches 0, it must stay there, since it is a nonnegative \mathbb{Q} -supermartingale. Therefore we have uniqueness in $[t, T]$. \square

By Propositions 5.1 and 5.3, and by Theorem 5.2, we get the following.

Corollary 5.4 $\widetilde{W} = V$ and it is the unique solution in \mathcal{C} to the HJB equation (8). Moreover, given $(t, x) \in [0, T] \times \mathbb{R}_+$, an optimal control in feedback form for (P) starting at (t, x) is given by

$$c_s^* = C^V(s, X_s^{t,x;V}), \quad \pi_s^* = \Pi^V(s, X_s^{t,x;V}), \quad (76)$$

where C^V, Π^V are the feedback maps defined in (69)-(70) associated to $V \in \mathcal{C}$, and where $X^{t,x;V}$ is the unique solution to (71) associated to C^V, Π^V .

Remark 5.5 As consequence of Proposition 5.1 and Corollary 5.4, we see that V satisfies the Inada condition $\frac{\partial}{\partial x} V(t, 0^+) = +\infty$ even if $\frac{\partial U_1}{\partial c}(\cdot, 0^+)$, $\frac{\partial U_1}{\partial x}(\cdot, 0^+)$ and $U_2'(0^+)$ (which are well defined by concavity) are all finite. Indeed, the fact that V satisfies the Inada condition at 0^+ is simply due to the fact that $x = 0$ is an absorbing boundary combined with Assumption 2.2(iv).

6 Applications

Current utility on the wealth may arise in several situations. For instance, we mention pension funds allocation (see, in a context of utility maximization, [8, 11] and, in a context of quadratic cost minimization, [9, 16]); optimal portfolio problems with random horizon (see [1, 4]); markets with illiquidity (see [12, 13]). We are going to describe the latter two applications.

6.1 Portfolio optimization with random horizon

A first application of our framework is to portfolio problems with random horizon. Consider the consumption/investment problem with state equation (1) when the time horizon of the agent is $T \wedge \tau$ where $T > 0$ is fixed and τ is some random variable $\tau \in [0, +\infty)$, i.e. the objective to maximize is a functional such as

$$\mathbb{E} \left[\int_0^{\tau \wedge T} G_1(t, c_t) dt + G_2(\tau \wedge T, X_{\tau \wedge T}) \right]. \quad (77)$$

In this context it is meaningful to assume, in general, that $\mathcal{F}_T \neq \mathcal{F}$, and that τ is just \mathcal{F} -measurable. A special case, which is the one we illustrate, as it may be covered by our framework, is when τ is independent of \mathcal{F}_T (this problem has been already treated in [1] in the case of terminal utility). Since τ is independent of $(\mathcal{F}_t)_{t \geq 0}$, setting $F(t) = \mathbb{P}\{\tau \leq t\}$ and assuming that F admits density f over $[0, T)$, the functional (77) may be rewritten as⁷

$$\mathbb{E} \left[\int_0^T (G_1(t, c_t)(1 - F(t)) + G_2(t, X_t)f(t)) dt + (1 - F(T))G_2(T, X_T) \right]. \quad (78)$$

So, it falls into our setting - under suitable assumptions on the functions G_1, G_2 - with

$$\begin{aligned} U_1(t, c, x) &= G_1(t, c)(1 - F(t)) + G_2(t, x)f(t), \\ U_2(x) &= (1 - F(T))G_2(T, x). \end{aligned}$$

⁷See [10] for the rewriting of the term corresponding to G_2 in the general case when τ may be dependent on \mathcal{F}_T , in which case one has to consider $F(t) := \mathbb{P}\{\tau \leq t \mid \mathcal{F}_t\}$.

Therefore we can apply our results, which allow to construct optimal feedback controls by Corollary 5.4. To this regard we notice that in [1] the regularity of the value function is *assumed* in the verification theorem, so the results given through the Dynamic Programming approach in [1] are definitively based on the possibility of finding (regular) explicit solutions to the HJB equation. Hence, while in [1] it is needed to take specific structures for the utility function, here we do not need that.

Finally, we observe that the rewriting of (77) as (78) can be performed also in the case $T = \infty$. So, applying our Remark 2.3(iv), we get that our results on the HJB equation and on the optimal feedback controls hold also in this case. The next subsection provides a significant example.

6.2 Market with liquid and illiquid assets

A related application of our results is the mixed liquid/illiquid investment model studied in [12, 13]. We refer to the latter references for details on the model.

Consider a market constituted by a riskless asset (assumed constant), and two risky assets L and I following Black-Scholes dynamics:

$$\begin{aligned} dL_t &= L_t (b_L dt + \sigma_L dW_t), & L_0 &= 1, \\ dI_t &= I_t (b_I dt + \sigma_I (\rho dW_t + \sqrt{1 - \rho^2} dB_t)), & I_0 &= 1, \end{aligned}$$

where W and B are independent Brownian motions, and $\rho \in (-1, 1)$ is a correlation parameter.

The specificity of the model is that, while the liquid asset L may be observed and traded continuously, the illiquid asset I may only be traded and observed at discrete random times $(\tau_k)_{k \geq 0}$, where we assume that $\tau_0 = 0$, and the interarrival times $\tau_{k+1} - \tau_k$ are i.i.d., and independent from (B, W) .

The investor's strategy is then a triple $((c_t)_{t \geq 0}, (\pi_t)_{t \geq 0}, (\alpha_k)_{k \in \mathbb{N}})$ where the components represent, respectively, the consumption, the amount invested in the liquid asset L at time t , and the amount invested in the illiquid asset I at time τ_k . The investor's wealth then follows the dynamics

$$\begin{aligned} R_0 &= r, \\ R_t &= R_{\tau_k} + \int_{\tau_k}^t (\pi_s (b_L ds + \sigma_L dW_s) - c_s ds) + \alpha_k \left(\frac{I_t}{I_{\tau_k}} - 1 \right), \quad t \in (\tau_k, \tau_{k+1}]. \end{aligned}$$

The investor aims at optimizing the following criterion

$$V(r) = \sup_{(c_t, \pi_t, \alpha_k) \in \mathcal{A}(r)} \mathbb{E} \int_0^\infty e^{-\beta s} U(c_s) ds,$$

where U is a utility function, the discount factor $\beta > 0$ is chosen large enough to guarantee finiteness to the problem, and the set $\mathcal{A}(r)$ is the set of admissible controls keeping the wealth nonnegative.

Let $\alpha_0 \in [0, r]$ and define, in the random interval $[0, \tau_1)$, the processes X, Y, J as

$$\begin{aligned} dX_t &= -c_t dt + \pi_t(b_L dt + \sigma_L dW_t), & X_0 &= r - \alpha_0, \\ dY_t &= Y_t \left(\frac{\rho b_L \sigma_I}{\sigma_L} dt + \rho \sigma_I dW_t \right), & Y_0 &= \alpha_0, \\ J_t &= \alpha_0 \frac{I_t}{Y_t}. \end{aligned}$$

In other words, X_t is the liquid wealth at time t (the wealth held in the riskless or in the liquid asset), $Y_t J_t$ is the wealth held in the illiquid asset I , and the total wealth is $R_t = X_t + Y_t J_t$.

We may apply a Dynamic Programming Principle between 0 and τ_1 , and see that V satisfies the following dynamic programming principle:

$$V(r) = \sup_{0 \leq \alpha_0 \leq r} \sup_{(c_t, \pi_t) \in \mathcal{A}'(r, \alpha_0)} \mathbb{E} \left[\int_0^{\tau_1} e^{-\beta s} U(c_s) ds + e^{-\beta \tau_1} V(R_{\tau_1}) \right], \quad (79)$$

where $\mathcal{A}'(r, \alpha_0)$ is the set of admissible controls (c_t, π_t) keeping the process X nonnegative in the interval $[0, \tau_1)$. Let us focus on the inner optimization problem in (79), i.e. assume that α_0 is fixed and we want to optimize only on $(c_t, \pi_t) \in \mathcal{A}'(r, \alpha_0)$, and let us show how this problem may be rewritten so as to fall in the framework of Subsection 6.1.

Let $\mathbb{F}^W = (\mathcal{F}^W)_{t \geq 0}$ denote the filtration generated by W . We note that Y is \mathbb{F}^W -adapted, while J is independent of \mathbb{F}^W . Moreover, since I is not observed in the interval $[0, \tau_1)$, the information available to the investor is given by the filtration \mathbb{F}^W in that interval. Hence, defining the function $(t, x, y) \mapsto G[V](t, x, y) := \mathbb{E}[V(x + y J_t)]$ and taking the conditional expectation with respect to $\mathcal{F}_{\tau_1}^W$ in the inner optimization problem of (79), this last one may be rewritten as

$$\sup_{(c_t, \pi_t) \in \mathcal{A}'(r, \alpha_0)} \mathbb{E} \left[\int_0^{\tau_1} e^{-\beta s} U(c_s) ds + e^{-\beta \tau_1} G[V](\tau_1, X_{\tau_1}, Y_{\tau_1}) \right]. \quad (80)$$

Now, if we choose $U(c) = \frac{c^p}{p}$, $p \in (0, 1)$, the value function V will be p -homogeneous, $V(r) = K_V \frac{r^p}{p}$, and we can reduce the state space of the above inner control problem to one space dimension. Indeed, let us consider the state variable $Z_t := \frac{X_t}{Y_t}$. Letting

$$\tilde{c}_s = \frac{c_s}{Y_s}, \quad \tilde{\theta}_s = \frac{\pi_s}{Y_s} - Z_s \frac{\rho \sigma_I}{\sigma_L},$$

one may check that Z is solution of the SDE

$$dZ_t = -\tilde{c}_t dt + \tilde{\theta}_t ((b_L - \rho \sigma_I \sigma_L) dt + \sigma_L dW_t), \quad Z_0 = z = \frac{r - \alpha_0}{\alpha_0}. \quad (81)$$

Furthermore, (80) may be rewritten as

$$\sup_{(\tilde{c}_t, \tilde{\theta}_t) \in \mathcal{A}''(z)} \mathbb{E} \left[\int_0^{\tau_1} e^{-\beta s} Y_s^p U(\tilde{c}_s) ds + e^{-\beta \tau_1} Y_{\tau_1}^p G[V](\tau_1, Z_{\tau_1}, 1) \right], \quad (82)$$

where $\mathcal{A}''(z)$ is the set of admissible controls $(\tilde{c}_t, \tilde{\theta}_t)$ keeping the process Z nonnegative. We may rewrite (82) just in terms of Z . In order to do that, notice that $Y_t^p = \alpha_0^p H_t e^{k_{Y,p} t}$, where

$k_{Y,p} = p\rho\frac{\sigma_I b_L}{\sigma_L} - \frac{p(1-p)\rho^2\sigma_I^2}{2}$ and H is a martingale defined by $H_0 = 1$, $dH_s = p\rho\sigma_I H_s dW_s$. Then, denoting by \mathbb{Q} the probability with density process H_t , we have that $\widehat{W}_t := W_t - p\rho\sigma_I t$ is a \mathbb{Q} -Brownian motion. Moreover, (81) is equivalent to

$$dZ_t = -\tilde{c}_t dt + \tilde{\theta}_t \left((b_L - \rho\sigma_I\sigma_L(1-p))dt + \sigma_L d\widehat{W}_t \right), \quad (83)$$

and the control problem can be rewritten as

$$\alpha_0^p \cdot \sup_{(\tilde{c}, \tilde{\theta}) \in \mathcal{A}''(z)} \mathbb{E}^{\mathbb{Q}} \left[\int_0^{\tau_1} e^{-(\beta - k_{Y,p})s} U(\tilde{c}_s) ds + e^{-(\beta - k_{Y,p})\tau_1} G[V](\tau_1, Z_{\tau_1}, 1) \right]. \quad (84)$$

Due to Subsection 6.1, the optimization problem (83)-(84) is now in the framework of this paper (as long as we assume that τ_1 has a density).

References

- [1] Blanchet-Scalliet C., El Karoui N., Jeanblanc M., and L. Martellini (2008), *Optimal Investment and Consumption Decisions when Time-Horizon is Uncertain*, Journal of Mathematical Economics, Vol. 44, No. 11, pp. 1100–1113.
- [2] Bian B., Miao S., and H. Zheng (2011), *Smooth Value Functions for a Class of Nonsmooth Utility Maximization Problems*, SIAM Journal on Financial Mathematics, Vol. 2, No. 1, pp. 727–747.
- [3] Bian B. and H. Zheng (2012), *Smooth Value Function with Applications in Wealth-CVaR Efficient Portfolio and Turnpike Property*. Arxiv preprint.
- [4] Bouchard B. and H. Pham (2004), *Wealth-path dependent utility maximization in incomplete markets*, Finance and Stochastics, Vol. 8, No. 4, pp. 579–603.
- [5] Choulli T., Taksar M., and X.Y. Zhou (2003), *A diffusion model for optimal dividend distribution for a company with constraints on risk control*, SIAM Journal on Control Optimization, Vol. 41, No. 6, pp. 1946–1979.
- [6] Crandall M., Ishii H., and P.L. Lions (1992), *User's Guide to Viscosity Solutions of Second Order Partial Differential Equation*, Bulletin of the American Mathematical Society, Vol. 27, pp. 1–67.
- [7] Cox J.C. and C. Huang (1989), *Optimal consumption and portfolio policies when asset prices follow a diffusion process*, Journal of Economic Theory, Vol. 49, pp. 33–83.
- [8] Di Giacinto M., Federico S., and F. Gozzi (2011), *Pension funds with minimum guarantee: a stochastic control approach*, Finance and Stochastics, Vol. 15, No. 2, pp. 297–342.
- [9] Di Giacinto M., Federico S., Gozzi F., and E. Vigna (2012), *Constrained portfolio choices in the decumulation phase of a pension plan*, Submitted, Preprint of SSRN.
- [10] Elliott R.J., Jeanblanc M., and M. Yor (2000), *Some models on default risk*, Mathematical Finance, Vol. 10, No. 2, pp. 179–196.

- [11] Federico S., *A stochastic control problem with delay arising in a pension fund model* (2011), Finance and Stochastics, Vol. 15, No. 3, pp. 421–459.
- [12] Federico S. and P. Gassiat (2012), *Viscosity characterization of the value function of an investment-consumption problem in presence of illiquid assets*, Submitted. Arxiv preprint.
- [13] Federico S., Gassiat P., and F. Gozzi (2012), *Impact of time illiquidity in a mixed market without full observation*. Submitted. Arxiv preprint.
- [14] Fleming W.H. and H.M. Soner (2006), *Controlled Markov Processes and Viscosity Solutions*, Second Edition, Springer-Verlag.
- [15] Friedman A. (1983), *Partial Differential Equations of Parabolic Type*, Reprint edition, Robert E. Krieger Publishing company.
- [16] Gerrard R., Haberman S., and E. Vigna (2004), *Optimal investment choices post-retirement in a defined contribution pension scheme*, Insurance: Mathematics and Economics, Vol. 35, No. 2, pp. 321–342
- [17] Kabanov Y. and M. Safarian (2009), *Markets with Transaction Costs: Mathematical Theory*, Springer-Verlag.
- [18] Kallenberg O. (2002), *Foundations of modern probability*, Second edition, Probability and its Applications, Springer-Verlag, New York.
- [19] Karatzas I., Lehoczky J., and S.E. Shreve (1987), *Optimal portfolio and consumption decisions for a “small investor” on a finite horizon*, SIAM Journal on Control and Optimization, Vol. 25, pp. 1557–1586.
- [20] Karatzas I. and S.E. Shreve (1998), *Methods of mathematical Finance*, Springer-Verlag, New York.
- [21] Krylov N.V. (1996), *Lectures on Elliptic and Parabolic Equations in Hölder Spaces*, Graduate Studies in Mathematics, Vol. 12, Ed. American Mathematical Society.
- [22] Lieberman G. (1996), *Second Order Partial Differential Equations*, World Scientific Press, Singapore.
- [23] Schwartz E. and C. Tebaldi (2006), *Illiquid assets and optimal portfolio choice*, NBER Working Paper No. w12633.
- [24] Rogers L.C.G. (2002), *Duality in constrained optimal investment and consumption problems: a synthesis*, Paris-Princeton Lectures on Mathematical Finance (Springer Lecture Notes in Mathematics 1814). Springer Berlin, pp. 95–131.
- [25] Yong J. and X.Y. Zhou (1999), *Stochastic Controls: Hamiltonian Systems and HJB Equations*, Springer Verlag, New York.
- [26] Zariphopoulou T. (1994), *Consumption-Investment Models with Constraints*, SIAM Journal of Control and Optimization, Vol. 32, No. 1, pp. 59–85.